



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Future developments

Citation for published version:

Telfer, E 2019, 'Future developments: *In vitro* growth (IVG) of human ovarian follicles', *Acta Obstetrica et Gynecologica Scandinavica*. <https://doi.org/10.1111/aogs.13592>

Digital Object Identifier (DOI):

[10.1111/aogs.13592](https://doi.org/10.1111/aogs.13592)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Acta Obstetrica et Gynecologica Scandinavica

Publisher Rights Statement:

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/aogs.13592

This article is protected by copyright. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 Future developments: *In vitro* growth (IVG) of human ovarian follicles

6 Evelyn E Telfer

8 Institute of Cell Biology and Genes and Development Group CDBS

9 The University of Edinburgh

10 The Hugh Robson Building

11 George Square

12 Edinburgh EH8 8XE

14 Disclosure:

15 Evelyn Telfer has no conflicts of interest to declare. Work relating to this area in the Telfer
16 laboratory has been funded by the Medical Research Council.

21 Key words: Oocyte/Follicle Culture/Primordial Follicles/IVG/IVA/IVM/

24 Key Message: In Vitro Growth (IVG) of immature oocytes from cryo-preserved ovarian
25 cortex has the potential to produce mature oocytes but is still at an experimental stage.

26 Further research is required before clinical application can be realised

Abstract

Removal and storage of ovarian cortical tissue is currently offered to young female cancer patients undergoing potentially sterilizing chemotherapy and/or radiotherapy. For patients at high risk of re-introduction of malignancy through auto-transplantation, the ultimate aim is to achieve complete oocyte development from this tissue *in vitro*. The ability to develop human oocytes from the earliest follicular stages through to maturation and fertilisation *in vitro* would revolutionise fertility preservation practice. This has been achieved in mouse where *in vitro* grown (IVG) oocytes from primordial follicles have resulted in the production of live offspring. Systems that support growth and development of oocytes from human ovarian cortex are being developed by several groups. This review focuses on the steps required to recapitulate *in vitro* the process of human oocyte development from the primordial stage and the systems currently available to support this.

Introduction

The ability to develop human immature oocytes *in vitro* would have many potential applications but would be of particular relevance to Fertility Preservation. Ovarian tissue cryopreservation is now an option for women with cancer prior to undergoing gonadotoxic treatments (1). Re-implantation of cryo-preserved tissue is currently the only option to use stored tissue but in many cases, particularly for women with Leukemia, malignant cells are present in the ovary therefore, re-implantation is not an option, however, the oocytes within this tissue could potentially be grown *in vitro* (1). Culture systems with the aim of achieving *in vitro* growth (IVG) of immature oocytes to maturity and subsequent fertilization *in vitro* (IVF) have been the subject of research for almost 40 years. Several systems that support the growth of later stages of follicle development from rodents have been developed (2-8) with some reporting the production of live young (3, 4, 6-8). Complete development from the most immature oocytes (primordial stages) *in vitro* with subsequent IVF of oocytes followed by embryo transfer and production of offspring has been achieved in the mouse using a two step culture system (9, 10). The initial studies resulted in the birth of one mouse which developed abnormalities as an adult (9). Following alterations to the culture medium several mouse embryos and offspring have been obtained using IVG oocytes then combined with *in vitro* maturation (IVM) and IVF (10). More recently, *in vitro* systems that support complete development of murine oocytes starting from induced pluripotent stem cells (11) and from primordial germ cells (12) have been reported and these systems have resulted in competent oocytes that produce embryos and live young.

The work on rodents has provided proof of principle and has encouraged the challenge of adapting these systems to support human oocyte development *in vitro*. IVG of primordial follicles would be of particular benefit to pre-pubertal girls undergoing fertility preservation before being exposed to damaging chemotherapy (13). Currently the only fertility preservation option for pre-pubertal girls is storage of ovarian cortical tissue with the potential for re-implantation at a later time (13). In cases where re-implantation is not possible, IVG could provide another option to restore fertility. For IVG to be clinically viable, the process would need to start with primordial follicles as cortical strips that are stored contain predominantly this stage. Whilst there still remains much detailed research to be carried out before IVG of primordial follicles could ever be clinically applied, a great deal of progress has been made in developing culture techniques to support human oocyte

development *in vitro*. In this review the current status of human IVG from primordial follicles will be considered.

Stages of Follicle Development

The majority of follicles within the ovary in all young mammalian females will be at the primordial stage of development and these follicles are continually utilised throughout reproductive life (14). It is not known whether the pool of primordial follicles represents a homogeneous population but at this stage follicles have not yet been exposed to selection processes that lead to follicle degeneration (15, 16). Primordial follicles represent a “resting” population of germ cells (oocyte arrested at dictyate stage of Prophase 1 of meiosis surrounded by a few flattened granulosa cells), that are formed pre-natally. Recruitment into growth takes place throughout the woman’s reproductive life and a sequence of precisely regulated processes is required for complete oocyte/follicle development to occur. The sequence starts with (a) initiation of primordial follicle growth and development to the preantral follicle stage; (b) growth of the preantral stage with formation of a fluid filled cavity (antrum) and expansion to the pre ovulatory or Graafian follicle stage (c) rupture of the Graafian follicle releasing a cumulus-oocyte complex at ovulation in response to the mid-cycle LH surge (17, 18).

The oocyte is held in meiotic arrest as it grows within the follicle and it must acquire the ability to resume meiosis (meiotic competence) as well as the ability to support fertilisation and embryonic development (developmental competence). Oocyte development is dependent upon the environment of the individual follicle for its function as a gamete and this is regulated by inhibitory and stimulatory endocrine, paracrine and autocrine signalling by the somatic cells of the follicle (granulosa and surrounding theca cells) enhanced by several oocyte specific factors mediated through bi-directional communication (19, 20). The physiological requirements of the oocyte, granulosa and theca cells are extremely complex and dynamic therefore recapitulating the process of follicle activation and growth *in vitro* is one of the greatest technical challenges in reproductive technology (18).

Growing Human follicles *in vitro*

Several approaches have been taken to support early human follicle development *in vitro* starting with primordial follicle activation (21-31). Whilst there are several culture systems that support a specific stage of human oocyte development *in vitro*, there is so far only one

report that supports human primordial follicles to the stage of meiotic maturation (30). To achieve complete development of human oocytes *in vitro* a multi-step culture system is required (26, 30, 32). The first step in this process is to facilitate the initiation of primordial follicle development and support early growth; the second step is to optimise the growth of follicles from preantral to antral stages; step 3 supports the completion of oocyte growth ready for *in vitro* maturation in step 4 (Figure 1). In optimising a culture system to obtain developmentally competent oocytes the focus should be on oocyte development and this may preclude the need to develop large follicular structures *in vitro*. The multi-step approach needs to support the changing requirements of the developing oocyte and its surrounding somatic (granulosa) cells whilst maintaining good oocyte-somatic cell interactions. Therefore, providing conditions that support the maintenance of appropriately differentiated somatic cells in contact with the developing oocyte similar to the oocyte granulosa cell complexes in the rodent system is essential (4, 10).

IVA: *In vitro* Activation of human primordial follicles

The majority of follicles within ovarian cortical tissue will be at the quiescent primordial stage. Activation of primordial follicles *in vitro* (IVA) and early follicle development are key features of any IVG system. Human primordial follicles can be activated and grow well within mechanically loosened cortical pieces, developing to multilaminar preantral (secondary) stages within 6 days (26, 30). Central to this process is preparation of the ovarian tissue. This involves removal of most of the underlying stromal tissue and any growing follicles so that the cultured tissue consists of ovarian cortex containing primordial and primary follicles. When these small fragments of human ovarian cortex are cultured there is a significant shift of follicles from the quiescent to the growing pool over short periods of 6 – 10 days (26, 29-31).

It remains unclear how follicle activation is controlled but it is known to involve a combination of inhibitory, stimulatory and maintenance factors (33). The importance of the phosphatidylinositol-3'-kinase (PI3K-AKT) signalling pathway within the oocyte has been implicated in regulating activation of primordial follicles using mouse knockout models (34) and in human using culture of ovarian cortex (29, 35, 36). The phosphatase and tensin homolog deleted on chromosome ten (PTEN) acts as a negative regulator of this pathway and suppresses initiation of follicle development (34). The transcription factor Forkhead Protein O3 (FOXO3) is a downstream effector of this pathway and acts to inhibit follicle recruitment

(37). Other components of this pathway are dependent on the mammalian target of rapamycin complex 1 (mTORC1), a serine/threonine kinase that regulates cell growth and proliferation in response to growth factors and nutrients and also regulates primordial follicle activation (38). From knockout mouse data it appears that whilst PTEN within the oocyte suppresses activation of primordial follicles mTORC promotes it. How these pathways regulate human follicle development is unclear but culture models facilitate the study of these processes (29, 36, 39).

Significant primordial follicle activation occurs in step one of the multi-step culture system (26, 30). This activation appears to be as a result of disrupting the Hippo signalling pathway during the preparation of the tissue (36, 40, 41). Hippo disruption increases expression of downstream growth factors but manipulation of the PI3K pathway results in further activation (29, 35, 36, 40, 41). Inhibition of PTEN in cultured human ovarian cortex results in increased activation of primordial follicles and more secondary follicles but subsequent growth and survival of isolated secondary follicles is compromised (29, 36). The detrimental effect on secondary follicles may be as a result of alterations in DNA damage and repair responses as demonstrated recently in a bovine culture model (42).

Cortical strip culture removes follicles from the *in vivo* endocrine and paracrine processes regulating growth rate; however, follicles will still be subject to the effect of follicle interactions and the influence of stromal cell factors. It is clear that tissue shape and stromal density are important factors that contribute to the regulation of follicle growth initiation *in vitro*, as solid cubes of cortical tissue show lesser growth initiation (21) than cortex cultured as flattened “sheets,” where much of the underlying stroma is removed (26, 30). The physical environment of the follicles within the cortical tissue affects their response to stimulatory and inhibitory factors and therefore influences their ability to grow (43). Once follicle growth has been initiated within cortical tissue they can develop to multi-laminar stages but do not survive well within the cortical environment as growth is inhibited resulting in loss of follicle integrity and oocyte survival (22, 26). Growing follicles need to be released from the cortical stromal environment and cultured individually to limit the effect of follicle interactions (26, 30, 31, 44).

IVG: *In vitro* Growth of human preantral follicles

173 Preantral follicles can be isolated from cortical tissue post culture by mechanical dissection,
174 enzymatic isolation or a combination of both. Collagenase and DNase can be used to remove
175 preantral follicles from stromal tissue, however collagenase can cause damage which leads to
176 poor survival of follicles (45). The presence of theca layers is required for growing follicles
177 to retain their structure and survive the second stage of IVG and these may be compromised
178 by collagenase (45). More purified enzyme preparations such as Liberase may avoid the
179 damage that occurs with Collagenase (46, 47). Mechanical isolation of follicles has the
180 advantage of preserving follicular integrity by maintaining the basal lamina and thecal layers,
181 however the procedure is laborious and results in a low yield (26, 30).

183 Supporting the growth of human preantral follicles *in vitro* has led to the development of
184 matrices to maintain follicle structure. Alginate hydrogels has been used to encapsulate
185 human preantral follicles and support their growth *in vitro* (48). Alginate encapsulation is
186 thought to mimic the extra cellular matrix *in vivo* in terms of its ability to facilitate molecular
187 exchange between the follicle and the culture medium whilst its flexibility can accommodate
188 cell proliferation but its rigidity prevents dissociation of the follicular unit. The rigidity of the
189 alginate capsule affects follicle development as inhibition of growth and reduced
190 steroidogenesis have been reported in murine follicles embedded in 1% alginate gels (49)
191 whereas fully grown human oocytes have been produced using 0.5% gels (48).

192 The application of tissue engineering to support the growth of isolated follicles has been
193 making progress; decellularized ovarian tissue and 3D micro-porous scaffolds are being
194 explored as matrices to support preantral follicle growth (50, 51). Recent work has explored
195 the production of electrospun patterned porous scaffolds which may be more accessible and
196 reproducible than decellularised tissue (52). Engineered scaffolds clearly have great potential
197 and should be developed further to support human follicle growth *in vitro*.

198 The multi-step culture system that has been developed for human follicles (30) does not use
199 matrices or scaffolds to support the growth of isolated preantral follicles. Isolated growing
200 follicles are cultured in v-shaped micro-well plates and this has supported follicular
201 architecture *in vitro* whilst promoting growth, differentiation and antral formation (26, 30).

203 Once follicles are isolated from the ovarian cortex their progression *in vitro* is remarkable.
204 Secondary human follicles isolated enzymatically from fresh ovarian tissue and cultured in
205 the presence of Follicle Stimulating Hormone (FSH), become steroidogenically active and

complete oocyte growth within 30 days (48) and these oocytes have been shown to be capable of meiotic maturation (53). Primordial follicles grown within fragments of ovarian cortex to multi-laminar stages which are then isolated without the use of enzymes and cultured in the presence of Activin and FSH become steroidogenic within 10 days of *in vitro* growth (26, 30).

During step 2 of the multi-step system, isolated follicles cultured individually form antral cavities within 6-8 days. At this stage oocyte-granulosa cell complexes can be retrieved by applying gentle pressure to the follicle. Complexes with complete cumulus and adherent mural granulosa cells are selected for step 3 of the multi-step system (Figure 1) (30). Step 3 involves culturing the complexes on membranes in the presence of Activin-A and rhFSH for up to 4 days until oocytes reach a diameter of 100 microns (30).

That *in vitro* grown follicles can produce fully grown oocytes after a relatively short culture period confirms that local ovarian factors inhibit follicle development *in vivo*. Oocyte size is an indicator of its ability to resume meiosis therefore sustaining oocyte growth is the major objective of any complete *in vitro* development system (17). There are clearly differences in growth rate depending on whether the whole follicle is cultured (48, 53) or whether, complexes are removed for further growth after an antral cavity has formed (26, 30).

Whether the growth rate observed *in vitro* should be characterised as accelerated is not clear. The rate observed represents uninhibited growth without brakes that are required *in vivo* to regulate follicle development within the context of the reproductive cycle. Comparisons of culture systems is needed to determine optimal conditions but at this time there is only one complete system that supports growth from human primordial to maturation (30). The next step is to determine whether the growth pattern *in vitro* can support the development of healthy oocytes or whether it is deleterious to oocyte function, epigenetic changes and health.

Meiotic Maturation of Oocytes from IVG human follicles

The final stage in the IVG process before IVF can take place is IVM to support resumption of meiosis to the point of Metaphase II (Figure 1). IVM has been a successful strategy for embryo production in domestic species and has been applied to human oocytes with varying degrees of success (54, 55). The first live birth after IVM of immature oocytes was reported in 1991 (56), although IVM was being utilised during the early development of IVF (57). IVM is performed in a limited number of centres and success rates vary with the main factor

being oocyte source and stage (55). The rate of maturation of immature oocytes remains below that of oocytes harvested from stimulated ovaries, indicating that the protocols are sub-optimal or many of the harvested oocytes are intrinsically unable to undergo maturation (54, 55).

IVG oocytes derived from the multistep culture system undergo meiotic maturation following an IVM protocol (30). These oocytes form Metaphase II spindles but emit abnormally large polar bodies (30). Polar body size is influenced by the proximity of the spindle to the oocyte cortex and the inter-chromosomal spacing within Metaphase II spindles (58). If there is a loss of spindle contact with the oocyte cortex this can lead to extrusion of large polar bodies (59). The cause of the large polar bodies in the IVG derived oocytes is not known but it indicates that culture conditions need to be further optimised. The consequences of such abnormalities on chromosome balance in mature oocytes needs to be investigated.

Whilst acknowledging that there are no fully optimised culture systems for human oocytes there is now proof of concept that complete *in vitro* growth of human oocytes is possible (30). The end point of any IVG system is to produce developmentally competent and epigenetically normal oocytes therefore future research needs to focus on optimising each of the stages and to gain further understanding of the epigenetic status of IVG oocytes and of any embryos formed (60).

Summary

The most significant clinical application of *in vitro* growth (IVG) of human oocytes is in Fertility Preservation given the widespread adoption of ovarian tissue cryopreservation for cancer patients (1). It is clear that making a good egg is not an easy or straightforward process (18). If reliable methodology could produce *in vitro* generated mature human oocytes capable of fertilisation this would be a viable alternative to autologous transplantation. Apart from the clinical implications and potential of the various *in vitro* growth systems; each of them provide access to the process of human oogenesis in an experimentally tractable paradigm. Through these systems we will gain greater understanding of human oocyte development which will ultimately lead to improvements in Fertility Preservation.

272

273

1. Anderson RA, Wallace WHB, Telfer EE. Ovarian tissue cryopreservation for fertility preservation: clinical and research perspectives. *Hum Reprod Open*. 2017; 2017:1-9.
2. Telfer E, Torrance C, Gosden RG. Morphological study of cultured preantral ovarian follicles of mice after transplantation under the kidney capsule. *J Reprod Fertil*. 1990;89:565-71.
3. Cortvrindt R, Smitz J, Van Steirteghem AC. In-vitro maturation, fertilization and embryo development of immature oocytes from early preantral follicles from prepuberal mice in a simplified culture system. *Hum Reprod*. 1996;11:2656-66.
4. Eppig JJ, Schroeder AC. Capacity of mouse oocytes from preantral follicles to undergo embryogenesis and development to live young after growth, maturation, and fertilization in vitro. *Biol Reprod*. 1989;41:268-76.
5. Jin SY, Lei L, Shikanov A, Shea LD, Woodruff TK. A novel two-step strategy for in vitro culture of early-stage ovarian follicles in the mouse. *Fertil Steril*. 2010;93:2633-9.
6. Roy SK, Greenwald GS. Hormonal requirements for the growth and differentiation of hamster preantral follicles in long-term culture. *J Reprod Fertil*. 1989;87:103-14.
7. Spears N, Boland NI, Murray AA, Gosden RG. Mouse oocytes derived from in vitro grown primary ovarian follicles are fertile. *Hum Reprod*. 1994;9:527-32.
8. Xu M, West E, Shea LD, Woodruff TK. Identification of a stage-specific permissive in vitro culture environment for follicle growth and oocyte development. *Biol Reprod*. 2006;75:916-23.
9. Eppig JJ, O'Brien MJ. Development in vitro of mouse oocytes from primordial follicles. *Biol Reprod*. 1996;54:197-207.
10. O'Brien MJ, Pendola JK, Eppig JJ. A revised protocol for in vitro development of mouse oocytes from primordial follicles dramatically improves their developmental competence. *Biol Reprod*. 2003;68:1682-6.
11. Hikabe O, Hamazaki N, Nagamatsu G, et al. Reconstitution in vitro of the entire cycle of the mouse female germ line. *Nature*. 2016;539:299-303.
12. Morohaku K, Tanimoto R, Sasaki K, et al. Complete in vitro generation of fertile oocytes from mouse primordial germ cells. *Proc Natl Acad Sci U S A*. 2016;113:9021-6.
13. Anderson RA, Mitchell RT, Kelsey TW, et al. Cancer treatment and gonadal function: experimental and established strategies for fertility preservation in children and young adults. *Lancet Diabetes Endocrinol*. 2015;3:556-67.
14. Gougeon A, Chainy GB. Morphometric studies of small follicles in ovaries of women at different ages. *J Reprod Fertil*. 1987;81:433-42.
15. Picton HM, Harris SE, Muruvi W, Chambers EL. The in vitro growth and maturation of follicles. *Reproduction*. 2008;136:703-15.
16. Thomas FH, Walters KA, Telfer EE. How to make a good oocyte: an update on in-vitro models to study follicle regulation. *Hum Reprod Update*. 2003;9:541-55.
17. Telfer EE, Zelinski MB. Ovarian follicle culture: advances and challenges for human and nonhuman primates. *Fertil Steril*. 2013;99:1523-33.
18. Anderson RA, Telfer EE. Being a good egg in the 21st century. *Br Med Bull*. 2018;127:83-9.
19. Li R, Albertini DF. The road to maturation: somatic cell interaction and self-organization of the mammalian oocyte. *Nat Rev Mol Cell Biol*. 2013;14:141-52.
20. Eppig JJ. Reproduction: Oocytes Call, Granulosa Cells Connect. *Curr Biol*. 2018;28:R354-R6.

21. Hovatta O, Silye R, Abir R, Krausz T, Winston RM. Extracellular matrix improves survival of both stored and fresh human primordial and primary ovarian follicles in long-term culture. *Hum Reprod.* 1997;12:1032-6.
22. Hovatta O, Wright C, Krausz T, Hardy K, Winston RM. Human primordial, primary and secondary ovarian follicles in long-term culture: effect of partial isolation. *Hum Reprod.* 1999;14:2519-24.
23. Wright CS, Hovatta O, Margara R, et al. Effects of follicle-stimulating hormone and serum substitution on the in-vitro growth of human ovarian follicles. *Hum Reprod.* 1999;14:1555-62.
24. Picton HM, Gosden RG. In vitro growth of human primordial follicles from frozen-banked ovarian tissue. *Mol Cell Endocrinol.* 2000;166:27-35.
25. Hreinsson JG, Scott JE, Rasmussen C, et al. Growth differentiation factor-9 promotes the growth, development, and survival of human ovarian follicles in organ culture. *J Clin Endocrinol Metab.* 2002;87:316-21.
26. Telfer EE, McLaughlin M, Ding C, Thong KJ. A two-step serum-free culture system supports development of human oocytes from primordial follicles in the presence of activin. *Hum Reprod.* 2008;23:1151-8.
27. Garor R, Abir R, Erman A, et al. Effects of basic fibroblast growth factor on in vitro development of human ovarian primordial follicles. *Fertil Steril.* 2009;91:1967-75.
28. Ding CC, Thong KJ, Krishna A, Telfer EE. Activin A inhibits activation of human primordial follicles in vitro. *J Assist Reprod Genet.* 2010;27:141-7.
29. McLaughlin M, Kinnell HL, Anderson RA, Telfer EE. Inhibition of phosphatase and tensin homologue (PTEN) in human ovary in vitro results in increased activation of primordial follicles but compromises development of growing follicles. *Mol Hum Reprod.* 2014;20:736-44.
30. McLaughlin M, Albertini DF, Wallace WHB, Anderson RA, Telfer EE. Metaphase II oocytes from human unilaminar follicles grown in a multi-step culture system. *Mol Hum Reprod.* 2018;24:135-42.
31. Anderson RA, McLaughlin M, Wallace WH, Albertini DF, Telfer EE. The immature human ovary shows loss of abnormal follicles and increasing follicle developmental competence through childhood and adolescence. *Hum Reprod.* 2014;29:97-106.
32. Smits J, Dolmans MM, Donnez J, et al. Current achievements and future research directions in ovarian tissue culture, in vitro follicle development and transplantation: implications for fertility preservation. *Hum Reprod Update.* 2010;16:395-414.
33. Nelson SM, Telfer EE, Anderson RA. The ageing ovary and uterus: new biological insights. *Hum Reprod Update.* 2013;19:67-83.
34. Reddy P, Liu L, Adhikari D, et al. Oocyte-specific deletion of Pten causes premature activation of the primordial follicle pool. *Science.* 2008;319:611-3.
35. Li J, Kawamura K, Cheng Y, et al. Activation of dormant ovarian follicles to generate mature eggs. *Proc Natl Acad Sci U S A.* 2010;107:10280-4.
36. Grosbois J, Demeestere I. Dynamics of PI3K and Hippo signaling pathways during in vitro human follicle activation. *Hum Reprod.* 2018;33:1705-14.
37. Castrillon DH, Miao L, Kollipara R, Horner JW, DePinho RA. Suppression of ovarian follicle activation in mice by the transcription factor Foxo3a. *Science.* 2003;301:215-8.
38. Adhikari D, Liu K. mTOR signaling in the control of activation of primordial follicles. *Cell Cycle.* 2010;9:1673-4.
39. McLaughlin M, Patrizio P, Kayisli U, et al. mTOR kinase inhibition results in oocyte loss characterized by empty follicles in human ovarian cortical strips cultured in vitro. *Fertil Steril.* 2011;96:1154-9.e1.

40. Hsueh AJ, Kawamura K, Cheng Y, Fauser BC. Intraovarian control of early folliculogenesis. *Endocr Rev.* 2015;36:1-24.
41. Kawamura K, Cheng Y, Suzuki N, et al. Hippo signaling disruption and Akt stimulation of ovarian follicles for infertility treatment. *Proc Natl Acad Sci U S A.* 2013;110:17474-9.
42. Maidarti M, Clarkson YL, McLaughlin M, Anderson RA, Telfer EE. Inhibition of PTEN activates bovine non-growing follicles in vitro but increases DNA damage and reduces DNA repair response. *Hum Reprod.* 2019;34:297-307.
43. McLaughlin EA, McIver SC. Awakening the oocyte: controlling primordial follicle development. *Reproduction.* 2009;137:1-11.
44. McLaughlin M, Telfer EE. Oocyte development in bovine primordial follicles is promoted by activin and FSH within a two-step serum-free culture system. *Reproduction.* 2010;139:971-8.
45. Telfer EE, Binnie JP, McCaffery FH, Campbell BK. In vitro development of oocytes from porcine and bovine primary follicles. *Mol Cell Endocrinol.* 2000;163:117-23.
46. Dolmans MM, Michaux N, Camboni A, et al. Evaluation of Liberase, a purified enzyme blend, for the isolation of human primordial and primary ovarian follicles. *Hum Reprod.* 2006;21:413-20.
47. Rice S, Ojha K, Mason H. Human ovarian biopsies as a viable source of pre-antral follicles. *Hum Reprod.* 2008;23:600-5.
48. Xu M, Barrett SL, West-Farrell E, et al. In vitro grown human ovarian follicles from cancer patients support oocyte growth. *Hum Reprod.* 2009;24:2531-40.
49. Heise M, Koepsel R, Russell AJ, McGee EA. Calcium alginate microencapsulation of ovarian follicles impacts FSH delivery and follicle morphology. *Reprod Biol Endocrinol.* 2005;3:47.
50. Laronda MM, Jakus AE, Whelan KA, et al. Initiation of puberty in mice following decellularized ovary transplant. *Biomaterials.* 2015;50:20-9.
51. Laronda MM, Rutz AL, Xiao S, et al. A bioprosthetic ovary created using 3D printed microporous scaffolds restores ovarian function in sterilized mice. *Nat Commun.* 2017;8:15261.
52. Liverani L, Raffel N, Fattahi A, et al. Electrospun patterned porous scaffolds for the support of ovarian follicles growth: a feasibility study. *Sci Rep.* 2019;9:1150.
53. Xiao S, Zhang J, Romero MM, et al. In vitro follicle growth supports human oocyte meiotic maturation. *Sci Rep.* 2015;5:17323.
54. Nogueira D, Sadeu JC, Montagut J. In vitro oocyte maturation: current status. *Semin Reprod Med.* 2012;30:199-213.
55. Chian RC, Uzelac PS, Nargund G. In vitro maturation of human immature oocytes for fertility preservation. *Fertil Steril.* 2013;99:1173-81.
56. Cha KY, Koo JJ, Ko JJ, et al. Pregnancy after in vitro fertilization of human follicular oocytes collected from nonstimulated cycles, their culture in vitro and their transfer in a donor oocyte program. *Fertil Steril.* 1991;55:109-13.
57. Edwards RG, Bavister BD, Steptoe PC. Early stages of fertilization in vitro of human oocytes matured in vitro. *Nature.* 1969;221:632-5.
58. Barrett SL, Albertini DF. Cumulus cell contact during oocyte maturation in mice regulates meiotic spindle positioning and enhances developmental competence. *J Assist Reprod Genet.* 2010;27:29-39.
59. Coticchio G, Guglielmo MC, Dal Canto M, et al. Mechanistic foundations of the metaphase II spindle of human oocytes matured in vivo and in vitro. *Hum Reprod.* 2013;28:3271-82.

60. Anckaert E, De Rycke M, Smits J. Culture of oocytes and risk of imprinting defects. Hum Reprod Update. 2013;19:52-66.

List of Abbreviations:

FSH	Follicle Stimulating Hormone
FOXO3	Forkhead box protein O3
IVA	In Vitro Activation
IVG	In Vitro Growth
IVF	In Vitro Fertilisation
IVM	In Vitro Maturation
PI3K-AKT	Phosphatidylinositol-3'-kinase and Protein Kinase B
PTEN	The phosphatase and tensin homolog deleted on chromosome ten
mTORC1	Mammalian target of rapamycin complex 1




Figure 1: Diagrammatic representation of a multi-step culture system to support in vitro growth (IVG) of oocytes from human primordial follicles through to maturation as described by McLaughlin et al., 2018. Step 1, in vitro activation within micro-cortex for 7 days (a) then micro-dissection of multi-laminar growing follicles to be placed in step 2 (b) and cultured individually until antral formation occurs (c) Step 3, Isolation of the oocyte granulosa complex (d) from the intact follicle for further growth on membranes for up to 4 days (e). Step 4, Oocyte-Cumulus complexes placed within medium for in vitro maturation (IVM). Oocytes are then analysed for the presence of a Metaphase II spindle and a polar body. Fertilisation of IVG human oocytes has not yet been tested.